

# Social Determinants of Health and Pediatric Long COVID in the US

Kyung E. Rhee, MD, MSc, MA; Tanayott Thaweethai, PhD; Deepti B. Pant, MPH; Cheryl R. Stein, PhD; Amy L. Salisbury, PhD; Patricia A. Kinser, PhD; Lawrence C. Kleinman, MD, MPH; Richard Gallagher, PhD; David Warburton, DSc, MD; Sindhu Mohandas, MD; Jessica N. Snowden, MD; Melissa S. Stockwell, MD, MPH; Kelan G. Tantisira, MD, MPH; Valerie J. Flaherman, MD, MPH; Ronald J. Teufel, MD, MSCR; Leah Castro, MA; Alicia Chung, EdD, MPH; Jocelyn Espinoza Esparza, MPH; Christine W. Hockett, PhD; Maria Isidoro-Chino, BA; Anita Krishnan, MD; Lacey A. McCormack, PhD, MPH; Aleisha M. Nabower, MD; Erica R. Nahin, PhD; Johana M. Rosas, PhD; Sarwat Siddiqui, BS; Jacqueline R. Szmuszkovicz, MD; Nita Vangeepuram, MD, MPH; Emily Zimmerman, PhD; Heather-Elizabeth Brown, MA; Megan Carmilani, MEd; K. Coombs, MTCM, Ac; Liza Fisher, MBA; Margot Gage Witvliet, PhD; John C. Wood, MD, PhD; Joshua D. Milner, MD; Erika B. Rosenzweig, MD; Katherine Irby, MD; Elizabeth W. Karlson, MD, MS; Zihan Qian, MS; Michelle F. Lamendola-Essel, DHSc, MS; Denise C. Hasson, MD; Stuart D. Katz, MD; H. Shonna Yin, MD, MSc; Andrea S. Foulkes, ScD; Rachel S. Gross, MD, MS; and the RECOVER-Pediatrics Group; and the ABCD-RECOVER Group; for the RECOVER-Pediatrics Consortium

## Supplemental content

**IMPORTANCE** Millions of children worldwide are experiencing prolonged symptoms after SARS-CoV-2 infection, yet social risk factors for developing long COVID are largely unknown. As child health is influenced by the environment in which they live and interact, adverse social determinants of health (SDOH) may contribute to the development of pediatric long COVID.

**OBJECTIVE** To identify whether adverse SDOH are associated with increased odds of long COVID in school-aged children and adolescents in the US.

**DESIGN, SETTING, AND PARTICIPANTS** This cross-sectional analysis of a multicenter, longitudinal, meta-cohort study encompassed 52 sites (health care and community settings) across the US. School-aged children (6-11 years; n = 903) and adolescents (12-17 years; n = 3681) with SARS-CoV-2 infection history were included. Those with an unknown date of first infection, history of multisystem inflammatory syndrome in children, or symptom surveys with less than 50% of questions completed were excluded. Participants were recruited via health care systems, long COVID clinics, fliers, websites, social media campaigns, radio, health fairs, community-based organizations, community health workers, and existing research cohorts from March 2022 to August 2024, and surveys were completed by caregivers between March 2022 and August 2024.

**EXPOSURE** Twenty-four individual social determinant of health factors were grouped into 5 Healthy People 2030 domains: economic stability, social and community context, caregiver education access and quality, neighborhood and built environment, and health care access and quality. Latent classes were created within each domain and used in regression models.

**MAIN OUTCOMES AND MEASURES** Presence of long COVID using caregiver-reported, symptom-based, age-specific research indices.

**RESULTS** The mean (SD) age among 4584 individuals included in this study was 14 (3) years, and 2330 (51%) of participants were male. The number of latent classes varied by domain; the reference group was the class with the least adversity. In unadjusted analyses, most classes in each domain were associated with higher odds of long COVID. After adjusting for many factors, including age group, sex, timing of infection, referral source, and other social determinant of health domains, economic instability characterized by difficulty covering expenses, poverty, receipt of government assistance, and food insecurity were associated with an increased risk of having long COVID (class 2 adjusted odds ratio [aOR], 1.57; 95% CI, 1.18-2.09; class 4 aOR, 2.39; 95% CI, 1.73-3.30); economic instability without food insecurity (class 3) was not (aOR, 0.93; 95% CI, 0.70-1.23). Poorer social and community context (eg, high levels of discrimination and low social support) was also associated with long COVID (aOR, 2.17; 95% CI, 1.77-2.66). Sensitivity analyses stratified by age group and adjusted for race and ethnicity did not alter or attenuate these results.

**CONCLUSIONS AND RELEVANCE** In this study, economic instability that included food insecurity and poor social and community context were associated with greater odds of pediatric long COVID. Those with food security, despite experiencing other economic challenges, did not have greater odds of long COVID. Further study is needed to determine if addressing SDOH factors can decrease the rate of pediatric long COVID.

JAMA Pediatr. doi:10.1001/jamapediatrics.2025.5485  
Published online January 5, 2026.

**Author Affiliations:** Author affiliations are listed at the end of this article.

**Group Information:** The RECOVER-Pediatrics Group and the ABCD-RECOVER Group appear at the end of the article. The RECOVER-Pediatrics Consortium members are listed in Supplement 2.

**Corresponding Author:** Kyung E. Rhee, MD, MSc, MA, Department of Pediatrics, Division of Child and Community Health, University of California, San Diego School of Medicine, Rady Children's Hospital San Diego, 9500 Gilman Dr, MC 0874, La Jolla, CA 92093 (k1rhee@health.ucsd.edu).

SARS-CoV-2 contributes to the development of long COVID, a postacute chronic condition lasting at least 3 months that can affect almost every organ system. Adolescents may experience loss of smell or taste, muscle or joint pain, fatigue, postexertional malaise, cognitive issues, and mood changes.<sup>1,2</sup> In younger children, abdominal pain and itchy skin are also reported. It is difficult to estimate the true prevalence of long COVID due to varying symptom clusters, potential relapsing or remitting course, and reliance on self-report data in children. Nevertheless, prospective studies have estimated that 10% of children have experienced long COVID,<sup>3-5</sup> translating to millions of youth worldwide.

The World Health Organization (WHO) defines social determinants of health (SDOH) as nonmedical factors that influence health outcomes and are related to the conditions in which one is born, grows, works, lives, and ages.<sup>6</sup> In the US, Healthy People 2030 characterized SDOH into 5 domains: (1) economic stability, (2) social and community context, (3) education access and quality, (4) neighborhood and built environment, and (5) health care access and quality.<sup>7</sup> Children who experience adverse SDOH have an increased risk of poor health outcomes, including the development of chronic diseases (eg, cardiovascular disease, type 2 diabetes, and cancer).<sup>8</sup> Adverse SDOH act as chronic stressors that activate the hypothalamic-pituitary-adrenal axis, triggering glucocorticoid production that modulates cardiovascular responses, metabolic activity, cognitive or emotional responses, and immune function.<sup>8-10</sup> Prolonged exposure can desensitize the hypothalamus, resulting in damage to multiple organs and influence the pathogenesis and progression of disease.<sup>10,11</sup> Chronic stressors, like poverty, food insecurity, and recurring discrimination, may increase inflammation and alter immune responses to toxic exposures, including infections.<sup>12</sup> Since long COVID is a postviral syndrome,<sup>13</sup> stress-related immune dysfunction may make it difficult for the body to combat persistent viral components, immune dysregulation, or tissue damage that result in long COVID symptoms.<sup>14</sup>

To date, most research examining the association between SDOH and COVID-related health outcomes has focused on the risk of contracting the SARS-CoV-2 virus. This evidence comes from ecological studies conducted during the first year of the pandemic and focuses on a few SDOH factors at a time. Generally, COVID-19 case rates were higher in locations with lower SDOH ratings (eg, unfavorable working, transportation, and living conditions,<sup>15-17</sup> fewer grocery stores,<sup>18</sup> air pollution,<sup>16,18</sup> lack of health insurance,<sup>17,18</sup> poor access to primary care,<sup>16,19</sup> lower COVID-19 vaccination rates,<sup>19</sup> and worse Social Vulnerability Index,<sup>20-22</sup> Area Deprivation Index,<sup>23</sup> and COVID-19 Community Vulnerability Index<sup>22</sup> scores). However, the association of adverse SDOH with the development of long COVID remains understudied.<sup>24</sup>

Few studies have examined the association between SDOH and prolonged COVID symptoms. These adult-focused studies show that lower economic and education status, housing insecurity or overcrowding, lack of social support, and discrimination may be associated with prolonged symptoms.<sup>25-30</sup> Similar research in children has been difficult due to variability in long COVID definitions and

## Key Points

**Question** What is the association between adverse social determinants of health and long COVID in school-aged children and adolescents?

**Findings** In this cross-sectional analysis of a meta-cohort study including 4584 US children and adolescents, households with greater economic instability and poorer social or community context (eg, low social support and high levels of discrimination) experienced significantly higher odds of pediatric long COVID. However, those experiencing food security, despite other economic challenges, did not have higher odds of long COVID; results were similar across age groups.

**Meaning** The findings of an association between adverse social determinants of health and greater likelihood of pediatric long COVID suggest that addressing adverse social drivers may mitigate future disease risk.

age-specific differences in symptoms.<sup>1</sup> Nevertheless, long COVID is especially concerning in pediatric populations because of the potential for long-term physical and mental health effects that could persist into adulthood.<sup>31</sup> Therefore, examining the association between adverse SDOH and pediatric long COVID is an important step toward understanding disease prevention.

The objective of this analysis was to examine the cross-sectional association between SDOH and long COVID in school-aged children and adolescents given that chronic stressors may alter immune functioning and increase one's susceptibility to postviral syndromes. Furthermore, since social factors rarely exist in isolation, there was a need to evaluate the combined effect of all SDOH domains to determine if certain domains were more salient than others. Therefore, using data from the pediatric National Institutes of Health-funded Researching COVID to Enhance Recovery (RECOVER) initiative, we examined the association between Healthy People 2030 SDOH domains and long COVID.

## Methods

### Study Design

The RECOVER Pediatric Observational Cohort Study (RECOVER-Pediatrics)<sup>1,32</sup> is one of the largest, nationally representative, combined retrospective and prospective, longitudinal, observational meta-cohort study of children and young adults (newborn to age 25 years old) and their caregivers focused on long COVID. Participants were recruited across the US from health care and community-based settings or from the extant National Institutes of Health-funded Adolescent Brain Cognitive Development (ABCD) Cohort study.<sup>33-35</sup> Cohort and recruitment strategies have been previously described.<sup>32</sup> RECOVER-Pediatrics is under the oversight of the New York University Grossman School of Medicine single institutional review board (de novo cohort) or the University of California San Diego Human Research Protections Program (ABCD cohort). Caregiver-child pairs completed written informed consent and an age-appropriate assent (aged 7

**Table 1. Demographic Characteristics of the Researching COVID to Enhance Recovery (RECOVER) Pediatric Observational Cohort, US, 2022 to 2024**

| Characteristic                                                        | No./total No. (%)                      |                                 |                    |
|-----------------------------------------------------------------------|----------------------------------------|---------------------------------|--------------------|
|                                                                       | School-aged children, 6-11 y (n = 903) | Adolescents, 12-17 y (n = 3681) | Overall (n = 4584) |
| Sex assigned at birth                                                 |                                        |                                 |                    |
| Female                                                                | 427/903 (47)                           | 1827/3681 (50)                  | 2254/4584 (49)     |
| Male                                                                  | 476/903 (53)                           | 1854/3681 (50)                  | 2330/4584 (51)     |
| Age at enrollment, mean (SD), y                                       | 9 (2)                                  | 15 (1)                          | 14 (3)             |
| Race and ethnicity <sup>a</sup>                                       |                                        |                                 |                    |
| American Indian or Alaska Native                                      | 24/895 (3)                             | 117/3667 (3)                    | 141/4562 (3)       |
| Asian                                                                 | 69/895 (8)                             | 236/3667 (6)                    | 305/4562 (7)       |
| Black or African American, non-Hispanic                               | 80/895 (9)                             | 388/3667 (11)                   | 468/4562 (10)      |
| Hispanic, Latino, or Spanish                                          | 332/895 (37)                           | 811/3667 (22)                   | 1143/4562 (25)     |
| Native Hawaiian or Other Pacific Islander                             | 1/895 (0)                              | 8/3667 (0)                      | 9/4562 (0)         |
| White, non-Hispanic                                                   | 381/895 (43)                           | 2073/3667 (57)                  | 2454/4562 (54)     |
| None of these fully describe me                                       | 8/895 (1)                              | 34/3667 (1)                     | 42/4562 (1)        |
| Missing                                                               | 8                                      | 14                              | 22                 |
| Recruitment or referral source                                        |                                        |                                 |                    |
| Existing, prospectively followed COVID cohort                         | 32/903 (4)                             | 10/3681 (0)                     | 42/4584 (1)        |
| Self-referral from RECOVER website or other unsolicited self-referral | 169/903 (19)                           | 221/3681 (6)                    | 390/4584 (9)       |
| Participant tested/treated in the health system                       | 214/903 (24)                           | 281/3681 (8)                    | 495/4584 (11)      |
| Community outreach                                                    | 267/903 (30)                           | 275/3681 (7)                    | 542/4584 (12)      |
| Existing non-COVID research or clinical cohort                        | 65/903 (7)                             | 2655/3681 (72)                  | 2720/4584 (59)     |
| Long COVID clinic                                                     | 20/903 (2)                             | 52/3681 (1)                     | 72/4584 (2)        |
| Community health center                                               | 75/903 (8)                             | 66/3681 (2)                     | 141/4584 (3)       |
| Public health department list                                         | 25/903 (3)                             | 68/3681 (2)                     | 93/4584 (2)        |
| Other                                                                 | 36/903 (4)                             | 53/3681 (1)                     | 89/4584 (2)        |
| Prevalent SARS-CoV-2 strain at first infection                        |                                        |                                 |                    |
| Pre-Omicron (before Dec 1, 2021)                                      | 479/903 (53)                           | 1949/3681 (53)                  | 2428/4584 (53)     |
| Omicron (Dec 1, 2021, or later)                                       | 424/903 (47)                           | 1732/3681 (47)                  | 2156/4584 (47)     |
| Vaccination status at first infection <sup>b</sup>                    |                                        |                                 |                    |
| Fully vaccinated                                                      | 220/903 (24)                           | 1547/3681 (42)                  | 1767/4584 (39)     |
| Partially vaccinated                                                  | 106/903 (12)                           | 266/3681 (7)                    | 372/4584 (8)       |
| Not eligible for vaccination                                          | 425/903 (47)                           | 1262/3681 (34)                  | 1687/4584 (37)     |
| Not vaccinated                                                        | 125/903 (14)                           | 484/3681 (13)                   | 609/4584 (13)      |
| Vaccinated but missing information                                    | 27/903 (3)                             | 122/3681 (3)                    | 149/4584 (3)       |
| Caregiver relationship to child                                       |                                        |                                 |                    |
| Mother                                                                | 807/871 (93)                           | 3289/3643 (90)                  | 4096/4514 (91)     |
| Someone else                                                          | 64/871 (7)                             | 354/3643 (10)                   | 418/4514 (9)       |
| Missing                                                               | 32                                     | 38                              | 70                 |
| Long COVID research index score                                       |                                        |                                 |                    |
| Mean (SD)                                                             | 3.1 (5.9)                              | 2.3 (4.8)                       | 2.4 (5.1)          |
| Median (IQR)                                                          | 0.0 (0.0-4.5)                          | 0.0 (0.0-2.0)                   | 0.0 (0.0-3.0)      |
| Long COVID status <sup>c</sup>                                        |                                        |                                 |                    |
| Long COVID-probable                                                   | 196/903 (22)                           | 580/3681 (16)                   | 776/4584 (17)      |
| Long COVID-unspecified                                                | 707/903 (78)                           | 3101/3681 (84)                  | 3808/4584 (83)     |

<sup>a</sup> Race and ethnicity were gathered via caregiver report and included in this analysis to enhance our understanding of how race and ethnicity may be associated with long COVID and social determinants of health. See the eMethods in Supplement 1 for additional details on how race and ethnicity were collected and categorized; note that participants could select more than 1 race and ethnicity group.

<sup>b</sup> See the eMethods in Supplement 2 for additional details on how vaccination categories were defined. Participants were ineligible for vaccination if their first infection date preceded when vaccines were available for children their age.

<sup>c</sup> Long COVID status is defined according to the long COVID research index as long COVID-probable if the long COVID research index score is 5.5 or greater for school-aged children or 5.0 or greater for adolescents. Otherwise, long COVID status is defined as long COVID-unspecified.

years and older). The study followed the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) reporting guideline.


### RECOVER-Pediatrics Participants

The analytic sample included children aged 6 to 17 years with a known SARS-CoV-2 infection and a baseline long COVID

symptom survey<sup>32</sup> completed by caregivers between March 21, 2022, and August 27, 2024. Baseline long COVID symptom surveys were completed at variable time points after the child's first SARS-CoV-2 infection. Surveys for prolonged symptoms must have been completed at least 90 days after the child's first infection but not within 30 days of a reinfection. Evidence of SARS-CoV-2 antibodies was not required. Surveys were avail-

**Figure 1. Proportion of Participants With Individual Social Determinants of Health Factors and Odds Ratios for Each Factor and Long COVID**

| Domain                                                         | Participants with factor, % | Among those with factor, % with long COVID | Among those without factor, % with long COVID | Total participants, No. | Missing, % | OR (95% CI)      | P value from $\chi^2$ test |
|----------------------------------------------------------------|-----------------------------|--------------------------------------------|-----------------------------------------------|-------------------------|------------|------------------|----------------------------|
| <b>Economic stability</b>                                      |                             |                                            |                                               |                         |            |                  |                            |
| Food insecurity                                                | 13.0                        | 31.5                                       | 13.3                                          | 3872                    | 15.5       | 3.00 (2.42-3.71) | <.001                      |
| Housing instability or homelessness                            | 2.8                         | 22.4                                       | 16.1                                          | 4088                    | 10.8       | 1.50 (0.96-2.35) | .09                        |
| Transportation difficulties                                    | 6.4                         | 30.2                                       | 15.4                                          | 4113                    | 10.3       | 2.38 (1.80-3.14) | <.001                      |
| Childcare difficulties                                         | 8.8                         | 26.4                                       | 15.3                                          | 4107                    | 10.4       | 1.99 (1.55-2.55) | <.001                      |
| Income <120% FPL                                               | 22.0                        | 23.9                                       | 14.1                                          | 3794                    | 17.2       | 1.91 (1.58-2.30) | <.001                      |
| Difficulty covering expenses                                   | 37.8                        | 24.2                                       | 11.7                                          | 3952                    | 13.8       | 2.40 (2.03-2.85) | <.001                      |
| Receipt of government financial assistance                     | 18.4                        | 27.6                                       | 13.8                                          | 4018                    | 12.3       | 2.38 (1.97-2.88) | <.001                      |
| Receipt of government nutrition support                        | 23.5                        | 25.5                                       | 13.6                                          | 4059                    | 11.5       | 2.19 (1.83-2.61) | <.001                      |
| <b>Social and community context</b>                            |                             |                                            |                                               |                         |            |                  |                            |
| Caregiver not living with partner or unmarried                 | 21.9                        | 22.2                                       | 14.3                                          | 4125                    | 10.0       | 1.70 (1.42-2.05) | <.001                      |
| Severe household overcrowding                                  | 6.5                         | 25.1                                       | 15.7                                          | 4107                    | 10.4       | 1.80 (1.35-2.40) | <.001                      |
| Childhood stressors                                            | 5.9                         | 32.0                                       | 14.4                                          | 3837                    | 16.3       | 2.79 (2.08-3.75) | <.001                      |
| Discrimination                                                 | 29.9                        | 24.8                                       | 12.9                                          | 4227                    | 7.8        | 2.23 (1.88-2.63) | <.001                      |
| Low social support                                             | 41.2                        | 22.8                                       | 12.9                                          | 4505                    | 1.7        | 2.00 (1.71-2.34) | <.001                      |
| Medical discrimination                                         | 3.4                         | 45.2                                       | 15.7                                          | 4496                    | 1.9        | 4.41 (3.18-6.12) | <.001                      |
| <b>Caregiver education access and quality</b>                  |                             |                                            |                                               |                         |            |                  |                            |
| Less than college education                                    | 36.9                        | 20.4                                       | 14.0                                          | 4121                    | 10.1       | 1.57 (1.33-1.86) | <.001                      |
| Low health literacy                                            | 2.1                         | 26.1                                       | 16.0                                          | 4095                    | 10.7       | 1.85 (1.14-3.00) | .02                        |
| <b>Neighborhood and built environment</b>                      |                             |                                            |                                               |                         |            |                  |                            |
| Lower neighborhood social cohesion                             | 21.6                        | 22.5                                       | 14.5                                          | 4087                    | 10.8       | 1.70 (1.41-2.05) | <.001                      |
| Disordered neighborhood                                        | 21.3                        | 21.3                                       | 14.9                                          | 4058                    | 11.5       | 1.54 (1.28-1.87) | <.001                      |
| Resides in a rural area                                        | 4.3                         | 25.6                                       | 16.5                                          | 4584                    | 0.0        | 1.74 (1.25-2.42) | .001                       |
| <b>Health care access and quality</b>                          |                             |                                            |                                               |                         |            |                  |                            |
| Child is uninsured or receives government health insurance     | 33.2                        | 21.6                                       | 14.6                                          | 4517                    | 1.5        | 1.62 (1.38-1.90) | <.001                      |
| Caregiver is uninsured or receives government health insurance | 38.2                        | 21.9                                       | 14.0                                          | 4497                    | 1.9        | 1.72 (1.47-2.01) | <.001                      |
| No health care visit in last 12 mo                             | 9.2                         | 8.2                                        | 17.7                                          | 4521                    | 1.4        | 0.41 (0.29-0.59) | <.001                      |
| Unmet medical needs                                            | 14.1                        | 35.6                                       | 13.8                                          | 4470                    | 2.5        | 3.45 (2.86-4.16) | <.001                      |
| Resides in a medically underserved area                        | 29.8                        | 18.8                                       | 16.1                                          | 4584                    | 0.0        | 1.20 (1.02-1.42) | .03                        |




Total participants refers to the number of participants who do not have missing data for that factor (of the total sample size of 4584). Participants with missing data are excluded from analyses presented in this figure. The odds ratios (ORs) and 95% CIs are shown for a logistic regression where long COVID is the outcome and the factor is the sole predictor. The P value from a  $\chi^2$  test for the association between each factor and long COVID is shown as well. FPL indicates federal poverty level.

able in English and Spanish. Participants with an unknown date of first infection, history of multisystem inflammatory syndrome in children, or symptom surveys with less than 50% of questions completed were excluded.

### Long COVID Outcome

The outcome was presence of long COVID at baseline based on RECOVER-Pediatrics age-specific long COVID research indices.<sup>1</sup> For this analysis, symptoms were considered prolonged if they

**Figure 2. Healthy People 2030 Social Determinants of Health Domain Latent Classes in the Researching COVID to Enhance Recovery (RECOVER) Pediatric Cohort, US, 2022-2024 (N = 4584)**



The radar plots and heatmaps show the proportion of individuals with each social determinant of health factor in each class. Class 1 reflects the class with the lowest average proportion of each social determinant of health factor in the study population. The displayed radar plots and heatmaps exclude participants with missing data (though missing data were accounted for when defining the latent classes via multiple imputation). FPL indicates federal poverty level.

were present at baseline, had persisted for at least 4 weeks, and were still present at the time of the survey (ie, at least 90 days after infection). For school-aged children (aged 6 to 11 years), long COVID-probable was defined as long COVID research index 5.5

or greater (range, 0.5-34.5) based on 10 symptoms (cognitive difficulties, back or neck pain, stomach pain, headache, fears about specific things, refusal to go to school, itchy skin or rashes, trouble sleeping, nausea or vomiting, and lightheadedness). For ado-

**Table 2. Estimated Odds Ratios for the Association Between Healthy People 2030 Social Determinants of Health (SDOH) Domain Latent Class and Long COVID**

| SDOH domain                            | Domain latent class <sup>a</sup> | No. (%)                   |                                           | OR (95% CI)             |                                                                                                    |                                                                                                                        |
|----------------------------------------|----------------------------------|---------------------------|-------------------------------------------|-------------------------|----------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------|
|                                        |                                  | Participants <sup>b</sup> | Participants with long COVID <sup>c</sup> | Unadjusted <sup>d</sup> | Adjusted for other SDOH domains, sex, age group, timing of infection, referral source <sup>e</sup> | Adjusted for other SDOH domains, sex, age group, timing of infection, referral source, race and ethnicity <sup>f</sup> |
| Economic stability                     | 1                                | 2975 (65)                 | 381 (12.8)                                | NA                      | NA                                                                                                 | NA                                                                                                                     |
|                                        | 2                                | 333 (7)                   | 77 (23.1)                                 | 1.75 (1.31-2.35)        | 1.57 (1.18-2.09)                                                                                   | 1.64 (1.23-2.18)                                                                                                       |
|                                        | 3                                | 657 (14)                  | 95 (14.5)                                 | 0.96 (0.67-1.38)        | 0.93 (0.70-1.23)                                                                                   | 0.98 (0.74-1.3)                                                                                                        |
|                                        | 4                                | 619 (14)                  | 223 (36.0)                                | 2.96 (2.23-3.92)        | 2.39 (1.73-3.30)                                                                                   | 2.52 (1.82-3.49)                                                                                                       |
| Social and community context           | 1                                | 2705 (59)                 | 351 (13.0)                                | NA                      | NA                                                                                                 | NA                                                                                                                     |
|                                        | 2                                | 1427 (31)                 | 246 (17.2)                                | 1.01 (0.79-1.30)        | 0.82 (0.64-1.05)                                                                                   | 0.84 (0.66-1.08)                                                                                                       |
|                                        | 3                                | 452 (10)                  | 179 (39.6)                                | 3.13 (2.60-3.77)        | 2.17 (1.77-2.66)                                                                                   | 2.18 (1.79-2.67)                                                                                                       |
| Caregiver education access and quality | 1                                | 2737 (60)                 | 388 (14.2)                                | NA                      | NA                                                                                                 | NA                                                                                                                     |
|                                        | 2                                | 1847 (40)                 | 388 (21.0)                                | 1.44 (1.19-1.74)        | 1.12 (0.96-1.3)                                                                                    | 1.18 (1.01-1.38)                                                                                                       |
| Neighborhood and built environment     | 1                                | 4018 (88)                 | 639 (15.9)                                | NA                      | NA                                                                                                 | NA                                                                                                                     |
|                                        | 2                                | 566 (12)                  | 137 (24.2)                                | 1.23 (0.96-1.57)        | 0.90 (0.73-1.12)                                                                                   | 0.94 (0.75-1.16)                                                                                                       |
| Health care access and quality         | 1                                | 3121 (68)                 | 441 (14.1)                                | NA                      | NA                                                                                                 | NA                                                                                                                     |
|                                        | 2                                | 1225 (27)                 | 262 (21.4)                                | 1.37 (1.05-1.80)        | 0.97 (0.76-1.23)                                                                                   | 1.00 (0.80-1.26)                                                                                                       |
|                                        | 3                                | 238 (5)                   | 73 (30.7)                                 | 2.09 (1.55-2.82)        | 1.14 (0.82-1.59)                                                                                   | 1.20 (0.86-1.66)                                                                                                       |

Abbreviations: NA, not applicable; OR, odds ratio.

<sup>a</sup> Participants are grouped into latent classes within each SDOH domain, meaning each participant belongs to 5 latent classes (1 for each domain). For example, a participant may belong to latent class 1 for economic stability, 2 for social and community context, 1 for caregiver education access and quality, 2 for neighborhood and built environment, and 3 for health care access and quality.

<sup>b</sup> The number of participants that belong to each latent class within each domain. These numbers sum to the total sample size within each domain. The denominator for each percentage is the total sample size (n = 4584).

<sup>c</sup> The number of participants within a given latent class with long COVID. The denominator for each percentage is the number of participants within the

given latent class.

<sup>d</sup> The estimated OR obtained from a logistic regression model with long COVID as the outcome and a single categorical predictor for the latent class within that domain. For example, this would be a 4-category variable for the economic stability domain. A separate model is fit for each domain.

<sup>e</sup> The estimated OR obtained from a logistic regression model with 5 categorical predictors, with 1 categorical variable representing the latent classes for each domain. The model is additionally adjusted for sex, age group, timing of infection, and recruitment or referral source. A single model is fit for all domains combined.

<sup>f</sup> The estimated OR from the same model as in the prior column, but with additional adjustment for race and ethnicity.

lescents (aged 12 to 17 years), long COVID-probable was defined as long COVID research index 5 or greater (range, 0.5-25) based on 8 symptoms (altered smell or taste, body or joint pain, fatigue, postexertional malaise, back or neck pain, cognitive difficulties, headache, and lightheadedness). All other participants were defined as long COVID-unspecified.

### Social Determinants of Health

RECOVER-Pediatrics collected data via 24 measures that align with the Healthy People 2030 SDOH domains.<sup>6</sup> SDOH were conceptualized through a multistep process. First, survey questions were mapped to individual SDOH factors. Second, individual SDOH factors were dichotomized to facilitate use in latent class analysis and enhance interpretability of results. Third, dichotomized factors were grouped into 5 SDOH domains. Detailed description of mapping and factor dichotomization is available in eTable 1 in [Supplement 1](#).

#### Economic Stability

Economic stability contained 8 factors: (1) food insecurity, assessed using the 10-item US Department of Agriculture Household Food Security module, defined as low or very low food security<sup>36</sup>; (2) housing instability, identified by living in transitional housing, staying in a shelter, or experiencing homelessness,<sup>37</sup> (3) transportation difficulties,

identified as difficulty getting to where they needed to go<sup>38</sup>; (4) childcare difficulties, identified as difficulty finding or keeping childcare<sup>37</sup>; (5) income less than 120% of the federal poverty level, based on state and household size-specific thresholds<sup>39</sup>; (6) difficulty covering expenses, including household expenses and bills; (7) receipt of government financial assistance, based on use of programs such as Temporary Assistance for Needy Families, Social Security, or Supplemental Social Security; and (8) receipt of government nutrition support, based on use of Supplemental Nutrition Assistance Program or Special Supplemental Nutrition Program for Women, Infants, and Children.

#### Social and Community Context

Social and community context contained 6 factors: (1) caregiver marital status, identified as being married or living with a partner vs not; (2) severe household overcrowding, defined according to WHO Housing and Health Guidelines (the number of people divided by the number of rooms in the household exceeds 1.5)<sup>40</sup>; (3) childhood stressors, assessed using a modified version of the 10-item Other Childhood Stressors scale, which assesses household relationship quality, bullying, incarceration, substance use, divorce, and neighborhood violence<sup>41</sup>; (4) discrimination experienced by the child, assessed using 9 items from the Everyday Discrimination Scale<sup>42</sup>;

(5) medical discrimination experienced by the child, assessed by 1 item from the Everyday Discrimination Scale (whether the caregiver believes their child is discriminated against while getting medical care); and (6) low social support, assessed using a shortened version of the Medical Outcomes Study Social Support Survey.<sup>43</sup>

#### Caregiver Education Access and Quality

Caregiver education access and quality contained 2 factors: (1) lower caregiver education, based on highest level of education completed (dichotomized as more or less than a bachelor's degree), and (2) low caregiver health literacy, measured using a validated 3-item screening tool which assessed needing help reading hospital materials, having problems learning about medical conditions, and low confidence in completing medical forms.<sup>44,45</sup>

#### Neighborhood and Built Environment

Neighborhood and built environment contained 3 factors: (1) lower neighborhood social cohesion (eg, helpfulness and trust in neighbors) using the 4-item Neighborhood Social Cohesion scale<sup>46</sup>; (2) disordered neighborhood, assessing elements such as neighborhood noise, vandalism, appearance, and abandoned buildings, using the 8-item Neighborhood Disorder scale<sup>47,48</sup>; and (3) living in a rural area as defined by the Health Resources and Services Administration.

#### Health Care Access and Quality

Health care access and quality contained 5 factors: (1) child does not have health insurance or receives health insurance from the government (eg, Medicaid and TRICARE); (2) caregiver does not have health insurance or receives health insurance from the government; (3) child did not have a health care visit in the past 12 months; (4) child has unmet health care needs (trouble getting needed health care visits or obtaining medical equipment or supplies); and (5) living in a medically underserved area as defined by the Health Resources and Services Administration.

#### Covariates

Demographic covariates included child sex, age, race, and ethnicity (eMethods in Supplement 1). Race and ethnicity data were collected via caregiver report and included in the study to enhance our understanding of how race and ethnicity may be associated with long COVID and SDOH. Survey covariates included recruitment or referral source and caregiver relationship to child. Clinical covariates included prevalent SARS-CoV-2 strain at time of first infection and vaccination status at first infection (eMethods in Supplement 1).

#### Statistical Analysis

First, we described the associations between each of the 24 dichotomized SDOH factors and long COVID using  $\chi^2$  tests and estimated unadjusted odds ratios using logistic regression. Second, we completed a latent class analysis using the dichotomized factors within each SDOH domain to address high levels of correlation across SDOH factors within domains.<sup>49,50</sup> As a result, each participant belonged to 1 latent class for each domain. This

approach allowed for equal weighting of each domain, rather than allowing the domain with the most factors (ie, economic stability) to be the primary driver of the identified latent classes. Missingness in SDOH was addressed using multiple imputation by chained equations with 25 imputations.<sup>51</sup> The procedure for identifying latent classes across imputations is described in the eMethods in Supplement 1. Across all domains, classes were ordered from lowest to highest average proportion across all factors. Therefore, the reference class (class 1) represented those with the fewest adverse SDOH factors, and the class with the highest number represented those with the most adverse SDOH factors. The caregiver education access and quality domain was not amenable to latent class analysis because it contained only 2 factors, so this domain was grouped a priori into 2 classes. Radar plots and heat maps were used to illustrate the prevalence of dichotomized SDOH factors within classes for each SDOH domain.

Third, we estimated whether the odds of long COVID differed across latent class domains in 3 sets of models. First, unadjusted models were fit for each latent class domain by fitting a logistic regression with a single categorical variable for latent class membership. Second, given the intersectionality of SDOH domains, we ran a combined model containing all 5 categorical variables (1 for each domain), adjusting for child age group, sex, time of infection (pre- vs post-Omicron, defined as first SARS-CoV-2 infection before vs on or after December 1, 2021), and recruitment or referral source (self-referral or community outreach vs other). Third, based on the National Academies of Sciences, Engineering, and Medicine's recommendation on using race and ethnicity identity in an analysis,<sup>22</sup> we repeated model 2 with an adjustment for child race and ethnicity. Comparing results with and without race and ethnicity could help determine how well our measures of SDOH captured the association of cultural identity with long COVID risk in children. To fit these models, generalized estimating equations were used with exchangeable correlation structure to account for clustering by recruitment site. Interpretation of results focused on estimated odds ratios where the lower bound of the 95% CI exceeds 1. Two-tailed P values less than .05 were considered statistically significant.

In sensitivity analysis, we repeated the latent class analysis and all regression analyses stratified by age group (school-aged children and adolescents). All statistical analyses were performed using the *poLCA*, *irr*, and *geepack* packages in R version 4.4.0.

## Results

After applying inclusion and exclusion criteria (eFigure 1 in Supplement 1), the study sample included 4584 participants (903 school-aged children and 3681 adolescents) recruited from 52 sites: 2330 (51%) of participants were male and 2254 (49%) were female, and the mean (SD) age was 14 (3) years (Table 1). As reported by the caregivers, 141 participants were American Indian or Alaska Native; 305 were Asian; 468 were Black or African American; 1143 were Hispanic, Latino, or Spanish; 9 were Native Hawaiian or Other Pacific Islander; 2454 were

White; 22 were missing race and ethnicity data; and 42 reported that none of the provided categories fully described them. More than half of participants (2428) were infected with SARS-CoV-2 before December 1, 2021 (pre-Omicron), and (776) 17% had long COVID based on the long COVID Research Index. The distribution of the long COVID research index is shown in eFigure 2 in Supplement 1.

Participants or their households who experienced adverse SDOH are presented in Figure 1. Most adverse SDOH factors across domains were associated with greater unadjusted odds of long COVID (Figure 1; eFigure 3 in Supplement 1). In the latent class analysis (Figure 2), 1658 participants (36%) did not experience any adverse SDOH in any domain, thus belonging in class 1 for all domains. Additionally, 1006 (22%) experienced adverse SDOH in only 1 domain, and 1920 (42%) experienced adverse SDOH in 2 or more domains. Of those experiencing adverse SDOH in 2 or more domains, 256 (13%) experienced adverse SDOH in all 5 domains. In each SDOH domain, individual classes were characterized by different SDOH factors. For example, in the economic stability domain, class 2 was predominantly characterized by difficulty covering expenses, class 3 by receipt of government nutrition support, and class 4 by food insecurity, income less than 120% of the federal poverty level, difficulty covering expenses, and receipt of government assistance. Model diagnostics for how the optimal numbers of latent classes were chosen are provided in eTable 2 in Supplement 1.

In each SDOH domain, the prevalence of long COVID was lowest in class 1 (the reference group) (Table 2). Across unadjusted models, the odds of long COVID were higher in economic stability class 2 and 4, social and community context class 3 (characterized by experiencing discrimination, low social support, and caregiver living alone), caregiver education access and quality class 2 (less than college and/or low health literacy), and health care access and quality class 2 (child and caregiver are uninsured or receive government health insurance) and class 3 (similar to class 2, plus living in a medically underserved area and having unmet medical needs).

Adjusting for sex, age group, time of infection, recruitment or referral source, and other SDOH domains attenuated all effect estimates, and the lower bound of the 95% CI for latent classes in the education and health care domains no longer exceeded 1. However, youth in economic stability class 2 still had higher odds of long COVID compared to youth in economic stability class 1 (adjusted odds ratio [aOR], 1.57; 95% CI, 1.18-2.09). Economic stability class 4 also had higher odds of long COVID (aOR, 2.39; 95% CI, 1.73-3.30). Class 3, characterized by economic instability without food insecurity, did not have higher odds of long COVID (aOR, 0.93; 95% CI, 0.70-1.23). Youth in social and community context class 3, predominantly characterized by experiencing discrimination and low social support, had 2.17 (95% CI, 1.77-2.66) times higher odds of long COVID compared to those in class 1. Additional adjustment for race and ethnicity did not substantively alter the pattern of results.

In sensitivity analyses, performing latent class analysis within age groups resulted in similar latent classes as identified in the primary analysis (not shown). Using the latent classes

identified in the primary analysis, stratification by age group only impacted the precision of the results due to smaller sample sizes (eTable 3 in Supplement 1).

## Discussion

In this cross-sectional analysis of a meta-cohort study, we found that US children living in households experiencing greater adversity in 2 SDOH domains, economic instability and adverse social and community context, were more likely to have pediatric long COVID compared to households experiencing the fewest social hardships. These associations were consistent between school-aged children and adolescents despite the different symptoms included in the age-specific long COVID research indices.<sup>1</sup> Multiple sensitivity analyses, including adjustment for race and ethnicity, did not meaningfully alter results. This is one of few studies that demonstrated an association between adverse SDOH and long COVID in children.<sup>24</sup> The breadth of data available in RECOVER-Pediatrics allowed for rich characterization of all 5 Healthy People 2030 SDOH domains and delineated the independent association of several SDOH with long COVID while accounting for the larger social context in which children were living.

In the US, it can be challenging to disentangle low socioeconomic status, discrimination, and other adverse SDOH factors from race and ethnicity because persons from racial and ethnic marginalized groups are more likely to be impacted by racism and stress related to their socioeconomic position.<sup>52</sup> There is some evidence that long COVID is more prevalent in Black and Hispanic adults<sup>53</sup> and that life stressors related to lower socioeconomic status may increase risk for long COVID.<sup>54</sup> In our analysis, we were able to demonstrate an association between long COVID and the social and community context domain, which incorporates measures of discrimination. This association did not change after adjustment for race and ethnicity, implying that discrimination and other social community factors were independently associated with long COVID.

The American Academy of Pediatrics recognized the association between adverse SDOH and child health in 2016 and reaffirmed it in 2021.<sup>55</sup> For children with long COVID, the combined effect of a serious viral infection and adverse social conditions may have increased their risk,<sup>56</sup> as chronic stress can lead to abnormal cortisol responses and immunologic derangements.<sup>57-59</sup> Thus, mitigating the potential impact of poverty and other adverse SDOH on child health is critical.<sup>55,60-63</sup> The COVID-19 pandemic bolstered some nutrition<sup>64</sup> and economic support programs,<sup>65-67</sup> and the expanded use of telehealth altered access to comprehensive health care. The impact of nutrition support programs may have been observed in our analysis as those experiencing economic instability without food insecurity were not at higher risk of long COVID, even after controlling for access to health care. It has been shown that some health behaviors, like exercise, sufficient sleep, and healthy diets, can decrease inflammation.<sup>68</sup> Efforts to support healthy lifestyle behaviors and strengthen an array of protective social factors particularly relevant to child health should be a policy priority.

## Limitations

This study has limitations. First, selection bias may have limited generalizability of these results. Families most affected by adverse SDOH could have been less likely to enroll, and study procedures were conducted in English and Spanish only. Second, SDOH data were reported by caregivers, and objective neighborhood-level data were unavailable. Third, since we used a research index of long COVID, we may have unintentionally miscategorized children with only 1 or 2 symptoms into the long COVID-unspecified group, resulting in conservative estimates of the associations between SDOH and long COVID. Furthermore, creation of latent classes in this analysis may not be similar to latent classes in other cohorts. Additionally, the factors in the study were dichotomized to facilitate latent class analysis and enhance interpretability. It is possible some conclusions may have been different if the variables were dichotomized in a different way, treated as continuous variables, or grouped into more categories. This was a cross-sectional investigation, which pre-

cludes causal inference and our understanding of the directionality of associations. Nevertheless, RECOVER-Pediatrics is poised to examine trajectories and identify long-term health impacts as the cohort continues to be followed.

## Conclusions

The results of this study demonstrate that long COVID is more prevalent among children living with economic instability and adverse social and community contexts. Given their young age and unclear trajectory of symptoms over the life course, the long-term economic effects and the health care needs of this population may be immense. Public health efforts to address modifiable adverse SDOH associated with long COVID are essential to potentially mitigate long COVID risk and promote child health more broadly as children continue to acquire COVID-19.

## ARTICLE INFORMATION

**Accepted for Publication:** November 5, 2025.

**Published Online:** January 5, 2026.

doi:10.1001/jamapediatrics.2025.5485

**Author Affiliations:** Department of Pediatrics, Division of Child and Community Health, University of California San Diego School of Medicine, Rady Children's Hospital, San Diego (Rhee); Massachusetts General Hospital Biostatistics, Boston (Thaweethai, Pant, Qian, Foulkes); Department of Medicine, Harvard Medical School, Boston, Massachusetts (Thaweethai, Foulkes); Department of Child and Adolescent Psychiatry, New York University Grossman School of Medicine, New York (Stein, Isidoro-Chino); School of Nursing, Virginia Commonwealth University, Richmond (Salisbury, Kinser, Espinoza Esparza); Department of Pediatrics, Division of Population Health, Quality, and Implementation Sciences, Rutgers Robert Wood Johnson Medical School, New Brunswick, New Jersey (Kleinman); Bristol Myers Squibb Children's Hospital, New Brunswick, New Jersey (Kleinman); Department of Child and Adolescent Psychiatry, Child Study Center, New York University Grossman School of Medicine, New York (Gallagher); Department of Pediatrics, Division of Neonatal-Perinatal Medicine, Children's Hospital Los Angeles, Keck School of Medicine, University of Southern California, Los Angeles (Warburton); Department of Pediatrics, Division of Infectious Diseases, Children's Hospital Los Angeles, Keck School of Medicine, University of Southern California, Los Angeles (Mohandas); Department of Pediatrics, Division of Infectious Diseases, University of Arkansas for Medical Sciences, Little Rock (Snowden); Department of Pediatrics, Division of Child and Adolescent Health, Columbia University Vagelos College of Physicians and Surgeons, New York, New York (Stockwell); Department of Population and Family Health, Mailman School of Public Health, New York-Presbyterian Hospital, New York (Stockwell); Department of Pediatrics, Division of Respiratory Medicine, University of California San Diego School of Medicine, Rady Children's Hospital, San Diego (Tantisira); Department of Pediatrics, Division of General Pediatrics, University of California, San Francisco (Flaherman); Department of Pediatrics,

Division of Pediatric Hospital Medicine, University of California San Diego School of Medicine, Rady Children's Hospital, San Diego (Teufel); Researching COVID to Enhance Recovery (RECOVER) Study, New York, New York (Castro, Brown, Carmilani, Coombs, Fisher, Witvliet); Department of Population Health, New York University Grossman School of Medicine, New York (Chung, Nahin, Rosas, Yin, Gross); Avera Research Institute, Sioux Falls, South Dakota (Hockett, McCormack); Department of Pediatrics, University of South Dakota School of Medicine, Sioux Falls (Hockett, McCormack); New York Medical College, Westchester Medical Center, Valhalla (Krishnan); Department of Pediatrics, University of Nebraska Medical Center and Children's Nebraska, Omaha (Nabower); David Geffen School of Medicine, University of California, Los Angeles (Siddiqui); Children's Hospital Los Angeles, Los Angeles, California (Szmuszkovicz); Icahn School of Medicine at Mount Sinai, New York, New York (Vangeepuram); Department of Communication Sciences & Disorders, Northeastern University, Boston, Massachusetts (Zimmerman); Long Covid Families, Charlotte, North Carolina (Carmilani); Vermont Center for Independent Living, Montpelier (Coombs); Department of Pediatrics, Division of Cardiology, Children's Hospital Los Angeles, Keck School of Medicine, University of Southern California, Los Angeles (Wood); Department of Pediatrics, Division of Pediatric Allergy, Immunology and Rheumatology, Columbia University Vagelos College of Physicians and Surgeons, New York, New York (Milner); Department of Pediatrics, Division of Cardiology, Columbia University Vagelos College of Physicians and Surgeons, New York, New York (Rosenzweig); Department of Pediatrics, Division of Critical Care, University of Arkansas for Medical Sciences, Little Rock (Irby); Department of Medicine, Division of Rheumatology, Inflammation, and Immunity, Massachusetts General Hospital, Boston (Karlson); Massachusetts General Brigham, Boston (Karlson); Department of Medicine, New York University Grossman School of Medicine, New York (Lamendola-Essel); Department of Pediatrics, Division of Pediatric Critical Care Medicine, New York University Grossman School of Medicine, New York (Hasson); Department of Medicine,

Division of Cardiology, New York University Grossman School of Medicine, New York (Katz); Department of Pediatrics, Division of General Pediatrics, New York University Grossman School of Medicine, New York (Yin, Gross); Department of Biostatistics, Harvard T. H. Chan School of Public Health, Boston, Massachusetts (Foulkes).

**The RECOVER-Pediatrics Group and the ABCD-RECOVER Group:** Judy L. Aschner, MD; Andrew M. Atz, MD; Dithi Banerjee, PhD; Amanda Bogie, MD; Hulya Bukulmez, MD; Katharine Clouser, MD; Lesley A. Cottrell, PhD; Kelly Cowan, MD; Viren A. D'Sa, MD; Allen J. Dozor, MD; Amy J. Elliott, PhD; E. Vince S. Faustino, MD, MHS; Alexander G. Fiks, MD, MSCE; Sunanda Gaur, MD; Maria L. Gennaro, MD; Stewart T. Gordon, MD, MPH; Uzma N. Hasan, MD; Christina M. Hester, PhD, MPH; Alexander H. Hogan, MD, MS; Daniel S. Hsia, MD; David C. Kaelber, MD, PhD, MPH; Jessica S. Kosut, MD; Sankaran Krishnan, MD, MPH; Russell J. McCulloh, MD; Ian C. Michelow, MD; Sheila M. Nolan, MD, MSCE; Carlos R. Oliveira, MD, PhD; Wilson D. Pace, MD, PhD; Paul Palumbo, MD; Hengameh Raissy, PharmD; Andy Reyes, MD; Judith L. Ross, MD; Juan C. Salazar, MD, MPH; Rangaraj Selvarangan, PhD; Michelle D. Stevenson, MD, MS; Alan Werzberger, MD; John M. Westfall, MD, MPH; Kathleen Zani, MD; William T. Zempsky, MD, MPH; James Chan, MA; Torri D. Metz, MD, MS; Jane W. Newburger, MD; Dongngan T. Truong, MD, MS; Candace H. Feldman, MD, MPH, ScD; Robin Auopperle, PhD; Fiona C. Baker, PhD; Marie T. Banich, PhD; Deanna M. Barch, PhD; Arielle Baskin-Sommers, PhD; James M. Bjork, PhD; Mirella Dapretto, PhD; Sandra A. Brown, PhD; B. J. Casey, PhD; Linda Chang, MD; Duncan B. Clark, MD, PhD; Anders M. Dale, PhD; Thomas M. Ernst, PhD; Damien A. Fair, PA-C, PhD; Sarah W. Feldstein Ewing, PhD; John J. Foxe, PhD; Edward G. Freedman, PhD; Naomi P. Friedman, PhD; Hugh Garavan, PhD; Dylan G. Gee, PhD; Raul Gonzalez, PhD; Kevin M. Gray, MD; Mary M. Heitzeg, PhD; Megan M. Herting, PhD; Joanna Jacobus, PhD; Angela R. Laird, PhD; Christine L. Larson, PhD; Krista M. Lisdahl, PhD; Monica Luciana, PhD; Beatriz Luna, PhD; Pamela A.F. Madden, PhD; Erin C. McGlade, PhD; Eva M. Müller-Dehring, PhD; Bonnie J. Nagel, PhD; Michael C. Neale, PhD; Martin P. Paulus, PhD; Alexandra S.

Potter, PhD; Perry F. Renshaw, MD, PhD; Elizabeth R. Sowell, PhD; Lindsay M. Squeglia, PhD; Lucina Q. Uddin, PhD; Sylvia Wilson, PhD; Deborah A. Yurgelun-Todd, PhD.

**Affiliations of The RECOVER-Pediatrics Group and the ABCD-RECOVER Group:** Department of Pediatrics, Division of Infectious Diseases, Children's Hospital Los Angeles, Keck School of Medicine, University of Southern California, Los Angeles (Ernst); Department of Pediatrics, Center for Discovery and Innovation, Hackensack Meridian School of Medicine, Nutley, New Jersey (Aschner); Medical University of South Carolina, Charleston, South Carolina (Atz); Children's Mercy Hospital and Clinics, Kansas City, Missouri (Banerjee); Department of Pediatrics, Oklahoma University Health Science Center, Oklahoma City (Bogie); MetroHealth System, Cleveland, Ohio (Bukulmez); Department of Pediatrics, Division of Pediatric Hospital Medicine, Hackensack Meridian School of Medicine, Nutley, New Jersey (Clouser, Reyes); Department of Pediatrics, West Virginia University, Morgantown (Cottrell); Department of Pediatrics, Division of Pediatric Pulmonology, University of Vermont, Burlington (Cowan); Department of Pediatrics, Rhode Island Hospital, Providence (D'Sa); Department of Pediatrics, Division of Pediatric Pulmonology, Allergy and Sleep Medicine, Boston Children's Health Physicians, New York Medical College, Valhalla (Dozor, Krishnan); Avera Research Institute, Sioux Falls, South Dakota (Elliott); Department of Pediatrics, Division of Critical Care, Yale University School of Medicine, New Haven, Connecticut (Faustino); Department of Pediatrics, Division of General Pediatrics, Children's Hospital of Philadelphia, Perelman School of Medicine at the University of Pennsylvania, Philadelphia (Fiks); Department of Pediatrics, Division of Allergy, Immunology, and Infectious Diseases, Rutgers Robert Wood Johnson Medical School, New Brunswick, New Jersey (Gaur); Department of Public Health, Medicine, Immunology, Microbiology, Infectious Diseases, Rutgers New Jersey Medical School, Newark (Gennaro); Pennington Biomedical Research Center, Clinical Trials Unit, Baton Rouge, Louisiana (Gordon, Hsia); Department of Pediatrics, Division of Infectious Diseases, Cooperman Barnabas Medical Center, Livingston, New Jersey (Hasan); DARTNet Institute, Aurora, Colorado (Hester, Pace, Westfall); Department of Pediatrics, Connecticut Children's, University of Connecticut School of Medicine, Hartford (Hogan, Zempsky); Departments of Pediatrics, Internal Medicine, and Population and Quantitative Health Sciences, MetroHealth System, Cleveland, Ohio (Kaelber); Department of Pediatrics, Division of Pediatric Hospitalist, University of Hawaii, John A Burns School of Medicine, Honolulu (Kosut); Department of Pediatrics, Division of Pediatric Hospital Medicine, University of Nebraska Medical Center, Omaha (McCulloh); Department of Pediatrics, Division of Infectious Diseases, Connecticut Children's, University of Connecticut School of Medicine, Hartford (Michelow, Salazar); Department of Pediatrics, Division of Infectious Diseases, Boston Children's Health Physicians, New York Medical College, Valhalla (Nolan); Department of Pediatrics, Division of Infectious Diseases, Yale University School of Medicine, New Haven, Connecticut (Oliveira); Department of Pediatrics and Medicine, Division of Infectious Disease and International

Health, Dartmouth-Hitchcock Medical Center, Lebanon, New Hampshire (Palumbo); Department of Pediatrics, Division of Pulmonary, University of New Mexico School of Medicine, Albuquerque (Raissy); Department of Pediatrics, Nemours Children's Health, Wilmington, Delaware (Ross); Department of Pediatrics, Children's Mercy Hospital and Clinics, Kansas City, Missouri (Selvarangan); Department of Pediatrics, Division of Norton Children's Emergency Medicine, University of Louisville School of Medicine, Louisville, Kentucky (Stevenson); Department of Pediatrics, Division of Child and Adolescent Health, Columbia University Vagelos College of Physicians and Surgeons, Best Healthcare, Monroe, New York (Werzberger); Department of Pediatrics, Division of Critical Care, University of Arkansas for Medical Sciences, Little Rock (Zani); Massachusetts General Hospital Biostatistics, Boston (Chan); Department of Obstetrics and Gynecology, Division of Maternal-Fetal Medicine, University of Utah Health, Salt Lake City (Metz); Department of Cardiology, Boston Children's Hospital; Department of Pediatrics, Harvard Medical School, Boston, Massachusetts (Newburger); Children's Healthcare of Atlanta Cardiology, Emory University School of Medicine, Atlanta, Georgia (Truong); Department of Medicine, Division of Rheumatology, Inflammation, and Immunity, Brigham and Women's Hospital, Boston, Massachusetts (Feldman); Department of Medicine, Harvard Medical School, Boston, Massachusetts (Feldman); Laureate Institute for Brain Research, Tulsa, Oklahoma (Aupperle, Paulus); Center for Health Sciences, Stanford Research Institute International, Menlo Park, California (Baker, Müller-Oehring); Institute of Cognitive Science and Department of Psychology and Neuroscience, University of Colorado Boulder, Boulder (Banich); Departments of Psychological & Brain Sciences, Psychiatry, and Radiology, Washington University in St Louis, St Louis, Missouri (Barch); Department of Psychology, Yale University, New Haven, Connecticut (Baskin-Sommers, Gee); Institute for Drug and Alcohol Studies, Virginia Commonwealth University, Richmond (Bjork); Department of Psychiatry and Biobehavioral Sciences, University of Southern California, Children's Hospital Los Angeles, Los Angeles (Dapretto, Uddin); Department of Psychology and Psychiatry, University of California San Diego School of Medicine, Rady Children's Hospital, San Diego (Brown); Department of Psychology, Barnard College, Columbia University, New York, New York (Casey); Department of Diagnostic Radiology and Nuclear Medicine, University of Maryland Baltimore, Baltimore (Chang); Department of Psychiatry, University of Pittsburgh, Pittsburgh, Pennsylvania (Clark, Luna); Departments of Neurosciences, Radiology, and Psychiatry, University of California San Diego School of Medicine, Rady Children's Hospital, San Diego (Dale); Institute of Child Development, Department of Pediatrics, Masonic Institute for the Developing Brain, University of Minnesota, Minneapolis (Fair); Oregon Health & Science University, Portland (Fair); Department of Psychology, University of Rhode Island, Kingston (Feldstein Ewing); Del Monte Institute for Neuroscience, Department of Neuroscience, University of Rochester School of Medicine, Rochester, New York (Foxe, Freedman); Institute for Behavioral Genetics and Department of Psychology and Neuroscience, University of Colorado, Boulder (Friedman); Department of

Psychiatry, University of Vermont, Burlington (Garavan); Department of Psychology, Florida International University, Miami (Gonzalez); Department of Psychiatry and Behavioral Sciences, Division of Addiction Sciences, Medical University of South Carolina, Charleston (Gray, Squeglia); Department of Psychiatry, University of Michigan, Ann Arbor (Heitzeg); Department of Population and Public Health Sciences, University of Southern California, Children's Hospital Los Angeles, Los Angeles (Herting); Department of Psychiatry, University of California San Diego School of Medicine, Rady Children's Hospital, San Diego (Jacobus); Department of Physics, Florida International University, Miami (Laird); Department of Psychology, University of Wisconsin, Milwaukee (Larson, Lisdahl); Department of Psychology, University of Minnesota, Minneapolis (Luciana); Department of Psychiatry, Washington University in St Louis, St Louis, Missouri (Madden); Department of Psychiatry, University of Utah School of Medicine, Salt Lake City (McGlade, Renshaw, Yurgelun-Todd); Rocky Mountain Mental Illness Research Education and Clinical Center, Department of Veterans Affairs, Salt Lake City Veterans Affairs Medical Center, Salt Lake City, Utah (McGlade, Renshaw, Yurgelun-Todd); Department of Psychiatry, Oregon Health & Science University, Portland (Nagel); Virginia Institute for Psychiatric & Behavioral Genetics, Virginia Commonwealth University, Richmond (Neale); Department of Psychiatry, Division of Clinical Neuroscience Research Unit, University of Vermont, Burlington (Potter); Department of Pediatrics, University of Southern California, Children's Hospital Los Angeles, Los Angeles (Sowell); Institute of Child Development, University of Minnesota, Minnesota (Wilson).

**Author Contributions:** Drs Thaweethai and Foulkes had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. Drs Rhee and Thaweethai are co-first authors. Drs Foulkes and Gross are co-senior authors.

**Concept and design:** Rhee, Thaweethai, Pant, Stein, Kinser, Kleinman, Gallagher, Warburton, Mohandas, Snowden, Teufel, Hockett, Nahin, Rosas, Szmuszkovicz, Vangeepuram, H. Brown, Carmilani, Fisher, Milner, Rosenzweig, Hasson, Katz, Foulkes, Gross, Bukulmez, Clouser, Cottrell, Kaelber, Kosut, Raissy, Salazar, Stevenson, Newburger, Feldman, S. Brown, Casey, Clark, Jacobus, Lisdahl, Luna, McGlade, Paulus, Squeglia.

**Acquisition, analysis, or interpretation of data:** Rhee, Thaweethai, Pant, Stein, Salisbury, Kleinman, Gallagher, Mohandas, Snowden, Stockwell, Tantisira, Flahearn, Castro, Chung, Espinoza Esparza, Hockett, Isidoro-Chino, A. Krishnan, McCormack, Nabower, Siddiqui, Szmuszkovicz, Vangeepuram, Zimmerman, Carmilani, Coombs, Fisher, Gage Witvliet, Wood, Rosenzweig, Irby, Karlson, Qian, Lamendola-Essel, Katz, Yin, Foulkes, Gross, Aschner, Atz, Banerjee, Bogie, Bukulmez, Cowan, D'Sa, Dozor, Elliott, Faustino, Fiks, Gaur, Gennaro, Gordon, Hasan, Hester, Hogan, Irby, Kaelber, S. Krishnan, McCulloh, Michelow, Nolan, Oliveira, Pace, Palumbo, Reyes Santos, Ross, Selvarangan, Stevenson, Werzberger, Westfall, Zani, Zempsky, Chan, Metz, Truong, Feldman, Aupperle, Baker, Banich, Barch, Baskin-Sommers, Bjork, Casey, Chang, Clark, Dale, Dapretto, Ernst, Fair, Feldstein Ewing, Foxe, Freedman, Friedman, Garavan, Gee, Gonzalez, Gray, Heitzeg, Herting,

Jacobus, Laird, Larson, Lisdahl, Luciana, Luna, Madden, McGlade, Müller-Oehring, NAGEL, Neale, Potter, Renshaw, Sowell, Uddin, Wilson, Yurgelun-Todd.

*Drafting of the manuscript:* Rhee, Thaweethai, Pant, Stein, Kinser, Gallagher, Teufel, Espinoza Esparza, Hockett, McCormack, Nahin, Vangeepuram, Gross, Bjork, Luna, Paulus.

*Critical review of the manuscript for important intellectual content:* Rhee, Thaweethai, Pant, Stein, Salisbury, Kleinman, Warburton, Mohandas, Snowden, Stockwell, Tantisira, Flaherman, Castro, Chung, Hockett, Isidoro-Chino, A. Krishnan, McCormack, Nabower, Rosas, Siddiqui, Szmuszkovicz, Vangeepuram, Zimmerman, H. Brown, Carmilani, Coombs, Fisher, Gage Witvliet, Wood, Milner, Rosenzweig, Irby, Karlson, Qian, Lamendola-Essel, Hasson, Katz, Yin, Foulkes, Gross, Aschner, Atz, Banerjee, Bogie, Bukulmez, Clouser, Cottrell, Cowan, D'Sa, Dozor, Elliott, Faustino, Fiks, Gaur, Gennaro, Gordon, Hasan, Hester, Hogan, Hsia, Kaelber, Kosut, S. Krishnan, McCulloh, Michelow, Nolan, Oliveira, Pace, Palumbo, Raissy, Reyes Santos, Ross, Salazar, Selvarangan, Stevenson, Werzberger, Westfall, Zani, Zempsky, Chan, Metz, Newburger, Truong, Feldman, Aupperle, Baker, Banich, Barch, Baskin-Sommers, Bjork, S. Brown, Casey, Chang, Clark, Dale, Dapretto, Ernst, Fair, Feldstein Ewing, Foxe, Freedman, Friedman, Garavan, Gee, Gonzalez, Gray, Heitzeg, Herting, Jacobus, Laird, Larson, Lisdahl, Luciana, Madden, McGlade, Müller-Oehring, NAGEL, Neale, Potter, Renshaw, Sowell, Squeglia, Uddin, Wilson, Yurgelun-Todd.

*Statistical analysis:* Thaweethai, Pant, Teufel, Vangeepuram, Qian, Foulkes, Chan, Neale.

*Obtained funding:* Rhee, Thaweethai, Salisbury, Kinser, Kleinman, Warburton, Snowden, Stockwell, Tantisira, Wood, Milner, Katz, Foulkes, Gross, Aschner, Cottrell, D'Sa, Elliott, Faustino, Raissy, Salazar, Selvarangan, Aupperle, Baker, Banich, Barch, Baskin-Sommers, Bjork, Casey, Chang, Clark, Dale, Ernst, Fair, Feldstein Ewing, Foxe, Friedman, Garavan, Gee, Gonzalez, Gray, Herting, Jacobus, Lisdahl, Luciana, Müller-Oehring, NAGEL, Renshaw, Sowell, Squeglia, Wilson.

*Administrative, technical, or material support:* Rhee, Stein, Snowden, Stockwell, Flaherman, Chung, Hockett, Isidoro-Chino, Nabower, Nahin, Rosas, Siddiqui, Zimmerman, Carmilani, Wood, Irby, Karlson, Qian, Lamendola-Essel, Hasson, Katz, Gross, Atz, Bukulmez, Dozor, Fiks, Hester, Hogan, Hsia, Gennaro, Gordon, Hogan, Hsia, Kaelber, S. Krishnan, McCulloh, Raissy, Ross, Stevenson, Westfall, Chan, Aupperle, Barch, Baskin-Sommers, Chang, Clark, Ernst, Feldstein Ewing, Foxe, Herting, Jacobus, Lisdahl, Luciana, Madden, McGlade, Paulus, Squeglia, Uddin.

*Supervision:* Rhee, Thaweethai, Kleinman, Mohandas, Snowden, Stockwell, Hockett, Nabower, Szmuszkovicz, H. Brown, Wood, Irby, Katz, Foulkes, Gross, Bukulmez, Dozor, Fiks, Hester, Hogan, Hsia, Kaelber, Kosut, S. Krishnan, Nolan, Oliveira, Raissy, Selvarangan, Feldman, Bjork, S. Brown, Chang, Dapretto, Ernst, Foxe, Garavan, Gee, Gonzalez, Heitzeg, Jacobus, Lisdahl, Madden, McGlade, Renshaw, Squeglia.

**Conflict of Interest Disclosures:** Dr Kleinman reported grants from the National Institutes of Health (NIH) during the conduct of the study; serving as a board member for Dartnet Institute, Anshe Emeth Memorial Temple, and Metuchen outside the submitted work; and stock ownership

in GSK, Regeneron, Sanofi, and Amgen. Dr Gallagher reported grants from the NIH during the conduct of the study. Dr Warburton reported grants from the NIH during the conduct of the study and outside the submitted work. Dr Snowden reported personal fees from Pfizer (COVID vaccine advisory board membership), now terminated, not related to this work, outside the submitted work. Dr Stockwell reported personal fees from RTI for administrative functions related to the NIH RECOVER Initiative; and a service agreement paid for role as Associate Director of the American Academy of Pediatrics' Pediatric Research in Office Settings to the trustees of Columbia University outside the submitted work. Dr Flaherman reported grants from the NIH and personal fees from Research Triangle International during the conduct of the study. Dr Teufel reported program support for trial from Sanofi, Enanta, and Moderna outside the submitted work. Dr Chung reported grants from the NIH during the conduct of the study. Dr Hockett reported grants from the National Heart, Lung, and Blood Institute (NHLBI) during the conduct of the study. Dr A. Krishnan reported grants from the NIH during the conduct of the study. Dr McCormack reported grants from the NIH during the conduct of the study. Dr Nabower reported grants from the NIH during the conduct of the study as well as personal fees from Aquifer (honorary for serving as associate editor) outside the submitted work. Dr Rosas reported grants from the NIH during the conduct of the study. Dr Szmuszkovicz reported grants from the NIH RECOVER project during the conduct of the study. Dr Carmilani reported serving as the volunteer president of Long Covid Families, a nonprofit organization that supports children affected by long COVID; Long Covid Families has received sponsorship funding from Moderna; Dr Carmilani did not receive any personal financial benefit from this sponsorship. Dr Wood reported grants from the NIH during the conduct of the study as well as personal fees from Agios (consulting), Bristol Myers Squibb (consulting), Hiilhurst Pharmaceuticals (data safety monitoring board and consulting), Regeneron (advisory board), Chiesi (educational session), and Pharmacosmos (consulting) outside the submitted work. Dr Milner reported grants from Pharming and personal fees from 1E Therapeutics, Amgen, and Sobi outside the submitted work. Dr Irby reported grants from the US Centers for Disease Control and Prevention (US Centers for Disease Control and Prevention) during the conduct of the study. Dr Lamendola-Essel reported grants from the NIH during the conduct of the study. Dr Katz reported grants from the NIH during the conduct of the study. Dr Foulkes reported grants from the NIH during the conduct of the study and personal fees from RoundTable outside the submitted work. Dr Gross reported grants from the NIH (to institution) during the conduct of the study and grants (to institution) from the NIH and the US Department of Agriculture outside the submitted work. Dr Aschner reported grants from the NIH during the conduct of the study. Dr Elliott reported grants from the NIH during the conduct of the study. Dr Faustino reported grants from the NIH during the conduct of the study. Dr Fiks reported salary support from the American Academy of Pediatrics for work as Pediatric Research in Office Setting Director, travel support for participation in the *JAMA Pediatrics* Editorial Board from the American Medical Association, honoraria for grand rounds from

Atlantic Health Systems and Children's Specialized Hospital, honoraria for Patient-Centered Outcomes Research Institute (PCORI) project advisory board from Boston Medical Center, honoraria for participation in an advisory panel related to the National Patient-Centered Clinical Research Network from Duke Clinical Research Institute, honoraria from Emory for Grand Rounds and another visit, grants from the NIH and PCORI, personal fees from Penn State (consulted on an NIH-funded project) and the University of California, Los Angeles (consultant on an NIH-funded project), honoraria for an invited lecture from the University of North Carolina at Chapel Hill, and research support to research team for a teen driving project from NJM Insurance outside the submitted work. Dr Hester reported grants from the NIH during the conduct of the study as well as grants from Exact Sciences, Genentech, DELFI Diagnostics, Abbott Diabetes Care, GSK, Amgen, Takeda, and PCORI outside the submitted work. Dr Kaelber reported grants from the NIH during the conduct of the study. Dr McCulloh reported grants from the NIH during the conduct of the study as well as personal fees from Pfizer for serving on a data monitoring committee and from Immutix for serving as member of the board of directors outside the submitted work. Dr Oliveira reported grants from the NIH during the conduct of the study. Dr Pace reported grants from the NIH during the conduct of the study as well as grants from Boehringer Ingelheim and AstraZeneca outside the submitted work. Dr Palumbo reported grants from the NIH during the conduct of the study. Dr Raissy reported grants from the NIH during the conduct of the study. Dr Reyes Santos reported grants from Hackensack Meridian Health during the conduct of the study. Dr Selvarangan reported grants from Biomerieux, Cepheid, Hologic, Meridian, Abbott, Diasorin, Qiagen, and Altona outside the submitted work. Dr Zani reported grants from the NIH during the conduct of the study. Dr Zempsky reported grants from the NIH during the conduct of the study as well as personal fees from Haleon and stock options from OmniHealth outside the submitted work. Dr Metz reported grants from the NIH during the conduct of the study as well as grants from Pfizer as a site principal investigator for a study of administration of Paxlovid in pregnancy for mild to moderate COVID-19 and grants from Moderna and Pfizer as a site principal investigator for a study of an RSV vaccine during pregnancy outside the submitted work. Dr Newburger reported grants from the NHLBI during the conduct of the study as well as grants from Pfizer and the NHLBI Pediatric Heart Network grant on sequelae of vaccine myocarditis (CAMP study) and grants from NHLBI (MUSIC study) on the multisystem inflammatory syndrome in children outside the submitted work. Dr Truong reported grants from the NIH during the conduct of the study as well as grants from Pfizer as co-principal investigator for a study on long-term follow-up of COVID-19 vaccine-associated myocarditis funded by Pfizer outside the submitted work. Dr Feldman reported grants from the NIH during the conduct of the study as well as grants from Bristol Myers Squibb (related to social determinants of health but unrelated to long COVID) outside the submitted work. Dr Barch reported grants from the National Institute of Mental Health and the National Institute on Drug Abuse during the conduct of the study. Dr Dale

reported other from Cortechs Labs (equity holder, director) outside the submitted work. Dr Ernst reported grants from the National Institute on Drug Abuse and the NIH during the conduct of the study as well as grants from the NIH and personal fees from the International Society for Magnetic Resonance in Medicine (Magnetic Resonance in Medicine deputy editor) outside the submitted work. Dr Fair reported other from Turing Medical (Founder and equity holder) outside the submitted work. Dr Friedman reported grants from the NIH during the conduct of the study. Dr Gray reported grants from Aelis Farma (site principal investigator of multisite trial) and consulting fees from Achieve Life Sciences and Indivior outside the submitted work. Dr Lisdahl reported grants from the NIH during the conduct of the study. Dr Luciana reported grants from the NIH to the University of Minnesota to fund data collection during the conduct of the study and for research outside the submitted work. Dr McGlade reported grants from the NIH during the conduct of the study. Dr Wilson reported grants from the NIH during the conduct of the study and outside the submitted work. No other disclosures were reported.

**Funding/Support:** This research was funded by National Institutes of Health agreements OT2HL161841, OT2HL161847, and OT2HL156812. Additional support came from grant R01 HL162373. The ABCD Study is supported by the National Institutes of Health and additional federal partners under award numbers U01DA041048, U01DA050989, U01DA051016, U01DA041022, U01DA051018, U01DA051037, U01DA050987, U01DA041174, U01DA041106, U01DA041117, U01DA041028, U01DA041134, U01DA050988, U01DA051039, U01DA041156, U01DA041025, U01DA041120, U01DA051038, U01DA041148, U01DA041093, U01DA041089, U24DA041123, U24DA041147. A full list of supporters is available at <https://abcdstudy.org/federalpartners.html>.

**Role of the Funder/Sponsor:** The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

**Group Information:** The RECOVER-Pediatrics Consortium members are listed in [Supplement 2](#).

**Disclaimer:** The content is solely the responsibility of the authors and does not necessarily represent the official views of the RECOVER Program, the National Institutes of Health, or other funders.

**Data Sharing Statement:** See [Supplement 3](#).

**Additional Contributions:** We would like to thank the National Community Engagement Group; all patient, caregiver, and community representatives; and all the participants enrolled in the RECOVER Initiative. Representatives are paid volunteers who receive compensation based on the hours worked.

## REFERENCES

- Gross RS, Thaweethai T, Kleinman LC, et al; RECOVER-Pediatrics Consortium; RECOVER-Pediatrics Group Authors. Characterizing Long COVID in children and adolescents. *JAMA*. 2024;332(14):1174-1188. doi:10.1001/jama.2024.12747
- Rao S, Gross RS, Mohandas S, et al. Postacute sequelae of SARS-CoV-2 in children. *Pediatrics*. 2024;153(3):e2023062570. doi:10.1542/peds.2023-062570
- Lopez-Leon S, Wegman-Ostrosky T, Ayuzo Del Valle NC, et al. Long-COVID in children and adolescents: a systematic review and meta-analyses. *Sci Rep*. 2022;12(1):9950. doi:10.1038/s41598-022-13495-5
- Pinto Pereira SM, Mensah A, Nugawela MD, et al; CLoCk Consortium. Long COVID in children and young after infection or reinfection with the Omicron variant: a prospective observational study. *J Pediatr*. 2023;259:113463. doi:10.1016/j.jpeds.2023.113463
- Behnood SA, Shafran R, Bennett SD, et al. Persistent symptoms following SARS-CoV-2 infection amongst children and young people: a meta-analysis of controlled and uncontrolled studies. *J Infect*. 2022;84(2):158-170. doi:10.1016/j.jinf.2021.11.011
- World Health Organization. Social determinants of health. Accessed November 17, 2024. [https://www.who.int/health-topics/social-determinants-of-health#tab=tab\\_1](https://www.who.int/health-topics/social-determinants-of-health#tab=tab_1)
- US Department of Health and Human Services, Office of Disease Prevention and Health Promotion. Healthy people 2030. Accessed January 21, 2025. <https://odphp.health.gov/healthypeople/objectives-and-data/social-determinants-health>
- Cockerham WC, Hamby BW, Oates GR. The social determinants of chronic disease. *Am J Prev Med*. 2017;52(1S1):S5-S12. doi:10.1016/j.amepre.2016.09.010
- Djuric Z, Bird CE, Furumoto-Dawson A, et al. Biomarkers of psychological stress in health disparities research. *Open Biomark J*. 2008;1:7-19. doi:10.2174/1875318300801010007
- Mariotti A. The effects of chronic stress on health: new insights into the molecular mechanisms of brain-body communication. *Future Sci OA*. 2015;1(3):FSO23. doi:10.4155/fso.15.21
- Silverman MN, Sternberg EM. Glucocorticoid regulation of inflammation and its functional correlates: from HPA axis to glucocorticoid receptor dysfunction. *Ann N Y Acad Sci*. 2012;1261:55-63. doi:10.1111/j.1749-6632.2012.06633.x
- Liu YZ, Wang YX, Jiang CL. Inflammation: the common pathway of stress-related diseases. *Front Hum Neurosci*. 2017;11:316. doi:10.3389/fnhum.2017.00316
- Choutka J, Jansari V, Hornig M, Iwasaki A. Unexplained post-acute infection syndromes. *Nat Med*. 2022;28(5):911-923. doi:10.1038/s41591-022-01810-6
- Davis HE, McCormell L, Vogel JM, Topol EJ. Long COVID: major findings, mechanisms and recommendations. *Nat Rev Microbiol*. 2023;21(3):133-146. doi:10.1038/s41579-022-00846-2
- Truong N, Asare AO. Assessing the effect of socio-economic features of low-income communities and COVID-19 related cases: an empirical study of New York City. *Glob Public Health*. 2021;16(1):1-16. doi:10.1080/17441692.2020.1850830
- Ali SMA, Sherman-Morris K, Meng Q, Ambinakudige S. Longitudinal disparities in social determinants of health and COVID-19 incidence and mortality in the United States from the three largest waves of the pandemic. *Spat Spatiotemporal Epidemiol*. 2023;46:100604. doi:10.1016/j.sste.2023.100604
- Sy KTL, Martinez ME, Rader B, White LF. Socioeconomic disparities in subway use and COVID-19 outcomes in New York City. *Am J Epidemiol*. 2021;190(7):1234-1242. doi:10.1093/aje/kwaa277
- De Jesus M, Ramachandra SS, Jafflin Z, et al. The environmental and social determinants of health matter in a pandemic: predictors of COVID-19 case and death rates in New York City. *Int J Environ Res Public Health*. 2021;18(16):8416. doi:10.3390/ijerph18168416
- Wylezinski LS, Harris CR, Heiser CN, Gray JD, Spurlock CF. Influence of social determinants of health and county vaccination rates on machine learning models to predict COVID-19 case growth in Tennessee. *BMJ Health Care Inform*. 2021;28(1):e100439. doi:10.1136/bmjhci-2021-100439
- Islam SJ, Nayak A, Hu Y, et al. Temporal trends in the association of social vulnerability and race/ethnicity with county-level COVID-19 incidence and outcomes in the USA: an ecological analysis. *BMJ Open*. 2021;11(7):e048086. doi:10.1136/bmjopen-2020-048086
- Lee J, Ramirez IJ. Geography of disparity: connecting COVID-19 vulnerability and social determinants of health in Colorado. *Behav Med*. 2022;48(2):72-84. doi:10.1080/08964289.2021.2021382
- McKinney J, Salmanian B, Grace R, et al. Social drivers of covid-19 disease severity in pregnant patients. *Am J Perinatol*. 2024;41(S01):e2269-e2278. doi:10.1055/a-2109-3876
- Khanna N, Klyushnenkova EN, Kaysin A. Association of COVID-19 with race and socioeconomic factors in family medicine. *J Am Board Fam Med*. 2021;34(suppl):S40-S47. doi:10.3122/jabfm.2021.51.200338
- Pellegrino R, Chiappini E, Licari A, Galli L, Marseglia GL. Prevalence and clinical presentation of long COVID in children: a systematic review. *Eur J Pediatr*. 2022;181(12):3995-4009. doi:10.1007/s00431-022-04600-x
- Lukkahatai N, Rodney T, Ling C, Daniel B, Han HR. Long COVID in the context of social determinants of health. *Front Public Health*. 2023;11:1098443. doi:10.3389/fpubh.2023.1098443
- Kogevinas M, Karachaliou M, Espinosa A, et al. Risk, determinants, and persistence of long-COVID in a population-based cohort study in Catalonia. *BMC Med*. 2025;23(1):140. doi:10.1186/s12916-025-03974-7
- Thomason ME, Hendrix CL, Werchan D, Brito NH. Perceived discrimination as a modifier of health, disease, and medicine: empirical data from the COVID-19 pandemic. *Transl Psychiatry*. 2022;12(1):284. doi:10.1038/s41398-022-02047-0
- Mukherjee S, Kshirsagar M, Becker N, et al. Identifying long-term effects of SARS-CoV-2 and their association with social determinants of health in a cohort of over one million COVID-19 survivors. *BMC Public Health*. 2022;22(1):2394. doi:10.1186/s12889-022-14806-1
- Packard SE, Susser E. Association of long COVID with housing insecurity in the United States, 2022-2023. *SSM Popul Health*. 2023;25:101586. doi:10.1016/j.ssmph.2023.101586
- Feldman CH, Santacroce L, Bassett IV, et al. Social determinants of health and risk for long COVID in the U.S. RECOVER-adult cohort. *Ann*

- Intern Med.* 2025;178(9):1287-1297. doi:10.7326/ANNALS-24-01971
31. Gupta M, Gupta N, Esang M. Long COVID in children and adolescents. *Prim Care Companion CNS Disord.* 2022;24(2):21r03218. doi:10.4088/PCC.21r03218
  32. Gross RS, Thaweethai T, Rosenzweig EB, et al; RECOVER-Pediatric Consortium. Researching COVID to enhance recovery (RECOVER) pediatric study protocol: rationale, objectives and design. *PLoS One.* 2024;19(5):e0285635. doi:10.1371/journal.pone.0285635
  33. Barch DM, Albaugh MD, Avenevoli S, et al. Demographic, physical and mental health assessments in the adolescent brain and cognitive development study: rationale and description. *Dev Cogn Neurosci.* 2018;32:55-66. doi:10.1016/j.dcn.2017.10.010
  34. Garavan H, Bartsch H, Conway K, et al. Recruiting the ABCD sample: design considerations and procedures. *Dev Cogn Neurosci.* 2018;32:16-22. doi:10.1016/j.dcn.2018.04.004
  35. Volkow ND, Koob GF, Croyle RT, et al. The conception of the ABCD study: from substance use to a broad NIH collaboration. *Dev Cogn Neurosci.* 2018;32:4-7. doi:10.1016/j.dcn.2017.10.002
  36. US Department of Agriculture. Measuring food security in the United States. guide to measuring household food security: revised 2000. Accessed November 18, 2025. <https://nhis.ipums.org/nhis/resources/FSGuide.pdf>
  37. National Institutes of Health. Rapid acceleration of Diagnostics (RADx) data hub. Accessed 11/18/2024. <https://radxdatahub.nih.gov/>
  38. Streuli S, Garfein RS, Gaines T, Fielding-Miller R. COVID-19 disproportionately impacts access to basic needs among households with disabled members. *Disabil Health J.* 2023;16(2):101443. doi:10.1016/j.dhjo.2023.101443
  39. American Council on Aging. 2025 Federal poverty levels: guidelines & how they determine Medicaid eligibility. Accessed November 11, 2024. <https://www.medicaidplanningassistance.org/federal-poverty-guidelines>
  40. World Health Organization. WHO Housing and Health Guidelines. Accessed November 18, 2025. <https://www.ncbi.nlm.nih.gov/books/NBK535293/>
  41. Sausen KA, Randolph JW, Casciato AN, Dietrich MS, Scholer SJ. The development, preliminary validation, and clinical application of the Quick Parenting Assessment. *Prev Sci.* 2022;23(2):306-320. doi:10.1007/s11211-021-01320-w
  42. Williams DR, Yan Y, Jackson JS, Anderson NB. Racial differences in physical and mental health: socio-economic status, stress and discrimination. *J Health Psychol.* 1997;2(3):335-351. doi:10.1177/135910539700200305
  43. Hays R, Sherbourne CD, Mazel R. *User's Manual for the Medical Outcomes Study (MOS) Core Measures of Health-Related Quality of Life.* RAND Corporation; 1994.
  44. Sarkar U, Schillinger D, López A, Sudore R. Validation of self-reported health literacy questions among diverse English and Spanish-speaking populations. *J Gen Intern Med.* 2011;26(3):265-271. doi:10.1007/s11606-010-1552-1
  45. Chew LD, Griffin JM, Partin MR, et al. Validation of screening questions for limited health literacy in a large VA outpatient population. *J Gen Intern Med.* 2008;23(5):561-566. doi:10.1007/s11606-008-0520-5
  46. Quinn TD, Wu F, Mody D, et al. Associations between neighborhood social cohesion and physical activity in the United States, National Health Interview Survey, 2017. *Prev Chronic Dis.* 2019;16:E163. doi:10.5888/pcd16.190085
  47. Elo I, Mykita L, Margolis R, Culhane J. Perceptions of neighborhood disorder: the role of individual and neighborhood characteristics. *Soc Sci Q.* 2009;90(5):1298-1320. doi:10.1111/j.1540-6237.2009.00657.x
  48. Ross CE, Mirowsky J. Neighborhood disorder, subjective alienation, and distress. *J Health Soc Behav.* 2009;50(1):49-64. doi:10.1177/002214650905000104
  49. McCarthy ML, Zheng Z, Wilder ME, et al. Latent class analysis to represent social determinant of health risk groups in the Medicaid cohort of the District of Columbia. *Med Care.* 2021;59(3):251-258. doi:10.1097/MLR.0000000000001468
  50. Weller B, Bowen N, Faubert S. Latent class analysis: a guide to best practice. *J Black Psychol.* 2020;46(4):287-311. doi:10.1177/0095798420930932
  51. White IR, Royston P, Wood AM. Multiple imputation using chained equations: Issues and guidance for practice. *Stat Med.* 2011;30(4):377-399. doi:10.1002/sim.4067
  52. US Centers for Disease Control and Prevention. Racism and health. Accessed November 17, 2024. <https://www.cdc.gov/minority-health/racism-health/index.html>
  53. Jacobs MM, Evans E, Ellis C. Racial, ethnic, and sex disparities in the incidence and cognitive symptomatology of long COVID-19. *J Natl Med Assoc.* 2023;115(2):233-243. doi:10.1016/j.jnma.2023.01.016
  54. Frontera JA, Sabadia S, Yang D, et al; NYU Neurology COVID-19 Study Team. Life stressors significantly impact long-term outcomes and post-acute symptoms 12-months after COVID-19 hospitalization. *J Neurol Sci.* 2022;443:120487. doi:10.1016/j.jns.2022.120487
  55. Schickedanz A, Szilagyi PG, Dreyer B. Child poverty and health in the United States: introduction and executive summary. *Acad Pediatr.* 2021;21(8S):S81-S85. doi:10.1016/j.acap.2021.09.003
  56. Boyce WT, Levitt P, Martinez FD, McEwen BS, Shonkoff JP. Genes, environments, and time: the biology of adversity and resilience. *Pediatrics.* 2021;147(2):e20201651. doi:10.1542/peds.2020-1651
  57. Shonkoff JP, Boyce WT, McEwen BS. Neuroscience, molecular biology, and the childhood roots of health disparities: building a new framework for health promotion and disease prevention. *JAMA.* 2009;301(21):2252-2259. doi:10.1001/jama.2009.754
  58. Alotiby A. Immunology of stress: a review article. *J Clin Med.* 2024;13(21):6394. doi:10.3390/jcm13216394
  59. Kasahara E, Inoue M. Cross-talk between HPA-axis-increased glucocorticoids and mitochondrial stress determines immune responses and clinical manifestations of patients with sepsis. *Redox Rep.* 2015;20(1):1-10. doi:10.1179/1351000214Y.0000000107
  60. Currie J, Chorniy A. Medicaid and child health insurance program improve child health and reduce poverty but face threats. *Acad Pediatr.* 2021;21(8S):S146-S153. doi:10.1016/j.acap.2021.01.009
  61. Empey A, Garcia A, Bell S. American Indian/Alaska Native child health and poverty. *Acad Pediatr.* 2021;21(8S):S134-S139. doi:10.1016/j.acap.2021.07.026
  62. Heard-Garris N, Boyd R, Kan K, Perez-Cardona L, Heard NJ, Johnson TJ. Structuring poverty: how racism shapes child poverty and child and adolescent health. *Acad Pediatr.* 2021;21(8S):S108-S116. doi:10.1016/j.acap.2021.05.026
  63. Jutte DP, Badruzzaman RA, Thomas-Squance R. Neighborhood poverty and child health: investing in communities to improve childhood opportunity and well-being. *Acad Pediatr.* 2021;21(8S):S184-S193. doi:10.1016/j.acap.2021.04.027
  64. Caspi C, Seligman H, Berge JM, Ng SW, Krieger J. COVID-19 pandemic era nutrition assistance: impact and sustainability. *Health Policy Brief.* Published online May 5, 2022. doi:10.1377/hpb20220330.534478
  65. US Economic Development Administration. CARES Act recovery assistance. Accessed November 26, 2025. <https://www.eda.gov/funding/programs/cares>
  66. US Economic Development Administration. American rescue plan: partnering with america's communities to build back better. Accessed November 26, 2025. <https://www.eda.gov/funding/programs/american-rescue-plan>
  67. Pandemic Response Accountability Committee. Where are they now? here's 7 pandemic relief programs that ended. Accessed November 26, 2025. <https://pandemicoversight.gov/news/articles/where-are-they-now-heres-7-pandemic-relief-programs-ended>
  68. Cuevas AG, Ong AD, Carvalho K, et al. Discrimination and systemic inflammation: a critical review and synthesis. *Brain Behav Immun.* 2020;89:465-479. doi:10.1016/j.bbi.2020.07.017